

***In vitro* investigation of the inhibitory effect of fucoidan on
 α -glucosidase enzyme activity**

Xin Yin

Bachelor of Pharmaceutical Engineering

Submitted in partial fulfillment of the requirements for the degree of

Master of Pharmaceutical Science

University of Tasmania

February 2013

Declaration of originality

This thesis contains no material that has been accepted for the award of any degree or diploma in any other tertiary institution.

To the best of knowledge and belief, this thesis contains no material that is previously published or written by any other except where due reference is made in the text of the thesis.

Xin Yin

February 2013

Authority of access

This thesis is not to be made available for loan or copying for one year following the date of this statement was signed. Following this time the thesis may be available for loan and limited copying in accordance with the *Copyright Act* 1968.

Xin Yin

February 2013

Table of Contents

Chapter 1 Literature Review

1.1 Marine algae and discovery history in Australia.....	2
1.2 The worldwide consumption, cultivation and utilisation of marine algae.....	2
1.3 Nutritional content and potential effects of marine algae.....	3
1.3.1 Protein and nitrogen.....	3
1.3.2 Fatty acid.....	4
1.3.3 Polysaccharide and dietary fibres.....	4
1.3.4 Minerals and vitamins.....	5
1.4 Fucoidan.....	5
1.4.1 General structure and physiochemical properties.....	5
1.4.2 structure–biological activities relationship.....	6
1.4.3 Different biological activities of the brown seaweed fucoidans.....	6
1.4.3.1 Anti–coagulant and anti–thrombotic activity.....	6
1.4.3.2 Anti–virals.....	7
1.4.3.3 Anti–tumor and anti–oxidant activities.....	7
1.4.3.4 Anti-inflammatory activity.....	7
1.4.4 Toxicity of fucoidan.....	8
1.5 Diabetes.....	8
1.5.1 Carbohydrate metabolism and postprandial hyperglycaemia.....	9
1.5.2 Currently available treatments of diabetes.....	10
1.5.2.1 Synthetic α –glucosidase inhibitors: acarbose and miglitol.....	10
1.5.2.2 Natural α –glucosidase inhibitors from plants and marine algae.....	11
1.6 Objectives.....	12
References.....	13

Chapter 2 Manuscript

2.1 Abstract.....	20
2.2 Introduction.....	21
2.3 Methodology.....	23
2.3.1 Materials.....	23
2.3.2 Mammalian α -glucosidase enzyme extraction.....	23
2.3.3 Determination of enzyme activity.....	23
2.3.4 Enzyme kinetics of α -glucosidase activity.....	24
2.3.5 Inhibitory assay.....	25
2.4 Results.....	27
2.4.1 Enzyme inhibitory assay of α -glucosidase.....	27
2.4.2 Enzyme activity of α -glucosidase.....	27
2.4.3 V_{\max} and K_m of α -glucosidase.....	27
2.5 Discussion.....	28
2.6 Conclusion.....	30
Declaration of interest.....	31
List of figures.....	32
References.....	34

List of figures

Figure 1 The inhibitory effect of fucoidan extracted from <i>Fucus synergy</i> , <i>Undaria pinnatifida</i> and <i>Fucus vesiculosus</i> on the activity of α -glucosidase, in comparison with acarbose at the same concentrations.....	33
Figure 2 Partial determination of enzyme kinetics of α -glucosidase using different concentrations of PNPG (ranging from 0.5 to 16 $\mu\text{g ml}^{-1}$).....	34

List of abbreviations

APTT	Activated partial thromboplastin time
CVDs	Cardiovascular diseases
DNA	Deoxyribonucleic acid
FAs	Fatty acids
IGT	Impaired dependent diabetes
K _m	Michaelis constant
LMWF	Low molecule weight fucoidan
mg	Milligram
ml	Milliliter
NF	Native fucoidan
NO	Nitric oxide
OSF	Oversulfated fucoidan
PNPG	4-Nitrophenyl β-D-glucuronide
PUFAs	Polyunsaturated fatty acids
RNA	Ribonucleic acid
STOP-NIDDM	Non-Insulin Dependent Diabetes Mellitus
V _{max}	Maximum reaction rate
µg	Microgram

Acknowledgment

First of all, I would like to express my appreciation to my supervisors, Professor Gregory Peterson (Head of School of Pharmacy, University of Tasmania), Dr. Rahul Patel (Masters in Pharmaceutical Science course coordinator), and Dr. Uttam Budhathoki for their suggestions and guidance and support thought out my research work, and help me submit this research thesis with success. Also, I wish to express my gratitude to Mrs Heather Galloway, Dr Peter Traill and Madhur Shastri for your generous assistance in the experiments.

Secondly, I want to thank my friends and parents for their selfless encouragement, support and love during my study.

Lastly but not the least, I am also thankful to Professor Gregory Peterson, Head of school of Pharmacy, who kindly provided me this opportunity to access this piece of research and the facilities of the department for the laboratory procedures.

Xin Yin

February 2013

Abstract

Introduction. Diabetes mellitus, a dysglycaemic disorder is causing a huge financial burden to society, government agencies and individuals every year. Diabetic patients are at the risk of long-term macro- and microvascular damages. Current therapies for diabetes include lifestyle modification and pharmacological interventions. However, conventional oral anti-hyperglycemic drugs produce a range of side effects such as hypoglycemia; therefore safer and more effective medications are required to manage diabetes and prevent the long-term complications. Fucooidan represents a large class of highly sulfated and water-soluble polysaccharides presents in all of the brown algae. Its antidiabetic effect has been reported in the literature. The objective of this study was to investigate the inhibitory effects of three different types of fucooidan on the activity of intestinal enzyme, α -glucosidase, which is involved in the facilitation of glucose absorption.

Methods. α -glucosidase enzyme was extracted from the rat-intestinal acetone powder. The enzyme activity of α -glucosidase was determined using an enzyme substrate 4-Nitrophenyl β -D-glucuronide (PNPG) and enzyme extract. The enzyme kinetics of α -glucosidase was performed using different concentrations of substrate. The calculation of V_{\max} and K_m were carried out based on Lineweaver-Burk plot. The inhibitory effect of three different types of fucooidan (*Undaria pinnatifida*, *Fucus vesiculosus*, *Fucus synergy*) was investigated. The stock solution containing 1000 $\mu\text{g ml}^{-1}$ of fucooidan was prepared. Different concentrations of fucooidan solutions were prepared by serially diluting the stock solution. Acarbose and Milli-Q water were used as positive control and negative control respectively. In enzyme assay, a reaction mixture contained enzyme extract, substrate and fucooidan.

Results and discussion. The activity of extracted α -glucosidase was found to be 0.0381 U ml^{-1} . The V_{\max} and K_m values were 0.2885 $\mu\text{g min}^{-1}$ and 8.8339 $\mu\text{g ml}^{-1}$ respectively. In the

inhibitory assay, the maximum percentage of inhibition of acarbose was 61.11%. In contrast, the maximum inhibitory effects of fucoidan extracted from *Fucus synergy* was 7.57%, followed by the extracts of *Fucus vesiculosus* (6.14%) and *Undaria pinnatifida* (3.30%).

Conclusion. Three different types of fucoidan investigated in the current study had negligible inhibitory effect on the activity of on α -glucosidase enzyme.

Chapter 1 Literature Review

1.1 Marine algae and discovery history in Australia

Marine algae, also known as seaweed is a wide group of organisms which require oxygen and light to produce nutrition for growth (1). Multiple taxonomical methods such as kingdom-division-class axis are developed to classify these organisms because it is impossible to place all of the existing algae into specific categories (2). However, the most common used category is based on their colour, namely phaeophytes (brown algae), rhodophytes (red algae), chlorophytes (green algae) (3) and cyanophytes (blue-green algae) (4). The extracts obtained from marine algae have various therapeutic functions, for instance, anti-tumor, antiviral and anti-coagulant activities (5).

The earliest description about Australian marine algae was made by a scientist from New South Wales as *Conferva umbilicata* in 1800 BC, which is currently known as *Microdictyon umbilicatum*. After that, Captain Vancouver collected marine algae systematically in his expedition in 1791. Thirty-nine years later, Charles Fraser became the first Australian who collected marine algae along the Swan River in Western Australia. Evidence has revealed that the Southern Australian region, including Tasmania is rich in marine algae resources, probably due to favorable growth conditions (6).

1.2 The worldwide consumption, cultivation and utilisation of marine algae

According to statistics reported in 2009, seaweeds contribute to approximately 25% of daily utilisation, including food consumption, agricultural field and health care industry (7). The annual consumption is dominated by a large portion of brown algae (66.5%), followed by figures of red algae (33%) and green algae (5%) (8). Interestingly, nearly 90% dry seaweed products just come from six countries, namely Chile, Korea, Japan, China, United Kingdom and France (9). Nevertheless, the consumption of seaweed products differs significantly from

Asia to Western countries. By comparison, seaweeds are not authorised as food in European countries until 1990 (10). With the epidemic requirement of seaweed products, artificial cultivation becomes the most productive way because of the high profits and ease of growth (11).

1.3 Nutritional content and potential effects of marine algae

Thirty years ago, the principal use of seaweeds was dominated by various industrial applications, for example, being used as gelling and thickening agents. Polysaccharide extracts such as alginate, carrageenan and agar did not used as food supplement until the year of 1990 in Europe. France was the first country to legislate them for human consumption (10). So far marine algae have been proved to be rich in various nutritional components such as vitamins, proteins, dietary fibres and minerals, and also have shown great interests in nutritional area. Undoubtedly, they are one of the promising food sources to cope with the shortage of food supplementation.

1.3.1 Protein and nitrogen

Generally, high protein and nitrogen content of marine algae is more important in food and agricultural industries (1). For example, some species show an increasing value for flavour formula because of high content of aspartate and glutamate (4). However, a species difference is observed for this aspect. Typically, brown seaweed contains relatively smaller proteinic percentage (average 5~15% of the dry weight), in comparison with that of red and green seaweed which are as high as 10~30%. Overall, marine algae are rich in proteins while poorer in digestibility (12). Although there are no substantial evidences indicating the digestibility of algal proteins, proteolytic enzymes such as pepsin and pancreatin are assumed to be responsible for the degradation of algal proteins (6,15,16).

1.3.2 Fatty acid

Lipid profiles of marine algae suggest that seaweeds have potential health benefit and turn out to be an option for hyperlipoidemic patients' diets (13). Seaweeds contain many essential fatty acids which are good for a balance diet, while low in lipid composition (between 1.5% and 4% of dry matter) (14). Marine algae have a remarkable level of polyunsaturated fatty acids (PUFAs) such as omega 3 (n-3) and omega 6 (n-6) fatty acids (FAs) and function well in preventing or modulating coronary heart disease. Some essential dietary fatty acids, for example, linoleic acids, are also detected in the seaweeds (15).

1.3.3 Polysaccharide and dietary fibres

Polysaccharides exist in many kinds of marine algae and express heterogeneous structures (16, 17). Algae polysaccharides can be divided into three major classes according to their functions and locations. The most common class is storage polysaccharides such as laminarin in brown seaweeds and starch in green seaweeds (16). Another common class is called cellular mucilage polysaccharides and other minor polysaccharides are found on the cell wall, for instance, carrageenans extracted from red seaweeds (14). Cell-wall polysaccharides have been proved to have various medicinal properties due to their sulfated groups (5, 20, 21). Most algae polysaccharides are proposed to be dietary fibre because there is a lack of digestive enzymes in human intestine; thus they cannot be considered as a reliable energy source (14). However, dietary fibre extracted from seaweeds contributes to the nutritional meal replacement for over-weight patients. Fucoidan, known as sulfated fucans, is an important member of sulfated polysaccharides and exhibit many important biological properties.

1.3.4 Minerals and vitamins

The mineral content of marine algae varies with growth environment, seasons, species and manufacturing methods. Generally, the composition of minerals is as high as 40%, containing various essential minerals, such as potassium, calcium, sodium, magnesium, phosphate, and a remarkable percentage of iodine (1, 12, 18). Therefore, edible marine algae have potential to be a prospective source of mineral supplementation product.

In addition to protein, dietary fibres and minerals, seaweeds are also considered to be a good source of vitamins. For example, red and green seaweeds contain significant levels of vitamin A and B (19), while part of the brown seaweeds are potential sources of vitamin C (12).

1.4 Fucoidan

1.4.1 General structure and physicochemical properties

Brown seaweeds produce multiple branched polysaccharides with species-specific sugar substituents. Algal fucoidans represent a family of highly sulfated and water-soluble polysaccharides presenting in all of the brown algae. They compose up to 25–30% of the seaweed dry weight. The biosynthesis pattern and the species diversity contribute to the heterogeneous structures of fucoidan (17, 20). Recent findings suggest that the backbone of fucoidan is constituted to α -(1 \rightarrow 3) linked fucose with substantial sulfate ester and L-fucose groups on C-2 or C-4 position (21). However, fucoidan structure is difficult to be analysed precisely due to the variability of linkage types and species deviation generally.

1.4.2 Structure–biological activities relationship

The algal fucans have some repeating pattern but are heterogeneous with the time of collection and species difference. It has been reported that their biological activities are

highly related with their structures. The position and amount of sulfated groups are two significant parameters that determine the biological activities of fucoidan (1, 22). It has also reported that molecule weight and size of fucoidan contribute to qualitative difference in previous biological studies (5, 25). For example, low molecule weight fucoidan (LMWF) (15-20 kDa) has a pro-angiogenic effect and better anti-coagulant outcomes than those in large size. Therefore, enzymatic degradation methods are used in the manufacturing process, aimed to produce a LMWF while retaining the maximum amount of sulfated groups (5).

1.4.3 Different biological activities of the brown seaweed fucoidans

One consequence of the heterogeneous structures of fucoidan is that each unique structure has a potential to be recognised by a specific receptor (5). Even though mechanisms by which fucoidan insert their biological actions remain controversial due to the complex structures, studies have indicated that fucoidan has considerably valuable therapeutic effects such as anti-coagulant (23, 24), anti-tumor (25), anti-inflammatory (16), antiviral (26, 27) and antioxidation (17).

1.4.3.1 Anti-coagulant and anti-thrombotic activity

Fucoidan has structural similarity in sulfated content with heparin, thus it exhibits anti-coagulant activity. A study that used activated partial thromboplastin time (APTT) mode has shown that fucoidan exhibits strong inhibitory effect on thrombin activity of platelet (28). Some studies illustrate that therapeutic outcomes and mode of action vary with the structure of fucoidan. For instance, increase in the sulfate content can improve the anti-coagulant activity by between 10 and 19% (21). Moreover, straight sulfated fucoidan show indirect inhibitory effect on thrombin while highly branched types can inhibit the activity of thrombin

without other factors (28). Because of the anti-coagulant activity of fucoidan, it can be considered as a promising alternative to heparin.

1.4.3.2 Anti-virals

In recent years, both *in vitro* and *in vivo* studies have demonstrated the antiviral activities of sulfated polysaccharides such as fucoidan. Fucoidan has an interesting function of anti RNA and DNA viruses (1). It has been reported that fucoidan could have protective antiviral activity via direct inhibition of viral replication and activating the host immune defense system (21).

1.4.3.3 Anti-tumor and anti-oxidant activities

Fucoidan is suggested to be a promising agent for anticancer and anti-tumor therapies. Both native and oversulfated fucoidans have been investigated. The results have shown that increasing the extent of sulfation can enhance the inhibition of tumor cell growth through stronger suppression on angiogenesis (29). Besides of its anti-tumor activity, fucoidans extracted from edible seaweeds are effective anti-oxidants.

1.4.3.4 Anti-inflammatory activity

The anti-inflammatory activity of fucoidan has also been investigated. Some extracts of fucoidan can decrease the neutrophil extravasation and transmigration at a dose of 15 mg/kg in an acute peritonitis rat model (28). The inhibitory effect of fucoidan on leukocyte recruitment in the inflammatory model may be associated with the release of nitric oxide (NO) (21).

1.4.4 Toxicity of fucoidan

Development and application of fucoidan as therapeutic agent has been an important research topic. However, data on its safety profile obtained from *in vitro* and *in vivo* studies of fucoidan are relatively limited. Two available research articles (30, 31) have reported the toxicities associated with fucoidan administration. Fucoidan extracted from *Undaria pinnatifida* did not cause genotoxicity when the oral dose was lower than 1000 mg/kg body weight per day in the tested animals. Whereas for fucoidan extracted from *Laminaria japonica* did not show signs of toxicity unless the dose was higher than 900 mg/kg body weight per day in the tested animals. The administration of fucoidan is considered to be safe when given at therapeutic doses. Nevertheless, the long-term side effects and toxicities associated with fucoidan require further investigation.

1.5 Diabetes

The incidents of diabetes mellitus, a dysglycaemic disorder is increasing world widely, causing a huge financial burden to society and individuals every year. According to the latest annual report of International Diabetes Federation, at the end of 2011, there are more than 366 million or 8.3 % of global adult population been diagnosed with diabetes, and the figure is predicted to be double at the end of 2030 (32). Diabetes is a metabolic disease characterized by high blood glucose level with the fasting glucose level more than 140 mg/dl (33). Diabetic patients are at the risk of long-term macro- and microvascular damages that can lead to further severe complications such as blindness, coronary artery disease and renal failure (34, 35).

Diabetes is classified into four main categories, namely type 1 or insulin-dependent diabetes, type 2 or non-insulin-dependent diabetes, gastrointestinal diabetes and other specific types

such as impaired glucose tolerance (IGT) (33, 35). Current antihyperglycaemic treatments only focus on controlling exacerbation and such available approaches have a range of adverse effects such as hypoglycemia, weight gain and nausea(36). Therefore, it is of importance to develop new antidiabetic therapies that could provide more efficient therapeutic outcomes with fewer side effects. In particular, decreasing the postprandial glucose levels through retarding the conversion of carbohydrates, fat and proteins into glucose is considered to be a promising approach because this method has a lower risk of hypoglycemia and weight gain which are frequently seen with the administration of other anti-diabetic agents. Nowadays, the inhibitory effects of natural products on digestive starch enzymes, for example, α -glucosidase, have drawn much attention in this aspect (37).

1.5.1 Carbohydrate metabolism and postprandial hyperglycaemia

Plasma glucose level is tightly controlled within a narrow range in nondiabetic people. After taking meals, increase in the postprandial glucose stimulates the secretion of insulin from pancreas β -cells (38). The balance between the rate of carbohydrate absorption from the gastrointestinal tract and the rate of utilisation is determined by several parameters such as nutrient type, digestive enzymes and reaction time of peripheral tissues to insulin secretion (39). In diabetic patients, postprandial hyperglycaemia could be observed as a consequence of insulin resistance or insulin deficiency (40). Evidences have suggested that postprandial hyperglycaemia itself is a significant risk factor of cardiovascular diseases (CVDs) and contributes to mortality rate (34). Studies have demonstrated that the risk of myocardial infarction is two to three fold greater in patients who have developed postprandial hyperglycaemia, compared with those with normal glucose tolerance (39). CVDs are associated with a soaring level of mortality in the patients with impaired dependent diabetes (IGT) and type 2 diabetes (34, 41).

1.5.2 Currently available treatments of diabetes

The main purpose of diabetic therapies is to control the blood glucose level and to prevent the complications such as CVDs (24, 29, 31). Lifestyle modification which involves dietary changes and exercise becomes a key strategy of prophylaxis. However, this approach alone is not sufficient for long-term glycaemic management. Therefore, pharmacological intervention is required to maintain the treatment outcomes (34). There are three main types of oral antidiabetics, namely insulin secretagogues, insulin sensitisers and inhibitors of carbohydrate absorption (38).

Recently, inhibitory effect of both synthesised and natural therapies on starch digesting enzymes such as α -glucosidase have been investigated since inhibiting the rate of carbohydrate absorption provides a safe way, with a lower risk of hypoglycaemia and greater benefits on preventing diabetic complications than conventional medications (36).

1.5.2.1 Synthetic α -glucosidase inhibitors: acarbose and miglitol

The development of α -glucosidase inhibitors such as acarbose and miglitol has provided a new approach in managing diabetes. Basically, these molecules reversibly bind to the carbohydrate region of α -glucosidase (an intestinal enzyme) This would reduce the digestion of carbohydrates and hence decrease the intestinal absorption of glucose (24, 28, 32). The major side effects of α -glucosidase inhibitors is gastrointestinal disturbance such as diarrhoea and abdominal discomfort, which are diminished with reduced dose (39). Acarbose and miglitol are two common drugs that can inhibit the activity of α -glucosidase enzyme. Diarrhoea, flatulence and abdominal distension are the major reported side effects of acarbose and miglitol. Large randomized clinical studies such as Non-Insulin Dependent Diabetes Mellitus (STOP-NIDDM) randomised trial (42, 43) and randomized double-blind

trial (44) have reported that both acarbose and miglitol showed superior outcomes in preventing postprandial hyperglycaemia and CVDs on patients with IGT or established type 2 diabetes, compared with insulin secretagogues and insulin sensitisers. Unlike the use of sulphonylureas and glinides, both of which are widely used for management of type 2 diabetes, administering acarbose or miglitol alone is not associated with weight gain and hypoglycaemia (34, 45). Additionally, acarbose and miglitol are safer to use with metformin and other oral antihyperglycaemic agents due to less drug-drug interactions (33, 35).

1.5.2.2 Natural α -glucosidase inhibitors from plants and marine algae

Other alternative sources for α -glucosidase inhibitors are plants and marine algae. The active extracts from plants and seaweeds provide more flexible food consumption choice for diabetic patients. Studies have shown that some plants extracts have hypoglycemic effects (37, 46, 47). For example, phenolic constituents of the heartwood of *Pterocarpus marsupium* can decrease the blood glucose level significantly in diabetic rat, in comparison of metformin (46). The anti-diabetic effect of both oversulfated fucoidan (OSF) and native fucoidan (NF) are also assessed via enzymatic reaction kinetics and their inhibitory activity against starch digestive enzymes (47). The results illustrated that the NF had little inhibitory effect on α -amylase while OSF can reduce the starch digestibility by 12%. Since fucoidan has a large molecular weight and low digestibility, it is less likely to be absorbed from the gastrointestinal tract. Hence, one possible mechanism of action for its potential anti-diabetic effects could be inhibition of intestinal enzymes.

1.6 Objectives

The aim of this study was to investigate the inhibitory effects of fucoindans from *Undaria pinnatifida*, *Fucus vesiculosus* and *Fucus synergy* on the activity of α -glucosidase.

References

1. KIM KTAE. Seasonal variation of seaweed components and novel biological function of fucoidan extracted from brown algae in Quebec. 2012.
2. Barsanti L, Gualtieri P. Algae: anatomy, biochemistry, and biotechnology: CRC; 2005.
3. Chapman VJ. Seaweeds and their uses: Taylor & Francis; 1950.
4. MacArtain P, Gill CIR, Brooks M, Campbell R, Rowland IR. Nutritional value of edible seaweeds. Nutrition reviews. 2007;65(12):535-43.
5. Morya V, Kim J, Kim EK. Algal fucoidan: structural and size-dependent bioactivities and their perspectives. Applied microbiology and biotechnology. 2012:1-12.
6. Womersley HBS. The marine algae of Australia. The Botanical Review. 1959;25(4):545-614.
7. Rao PS, Ganesan K, Kumar KS. Seaweeds: A Survey of Research and Utilization. Algal Biology and Biotechnology, Krishan Makhijanj, New Delhi. 2009.
8. Marinho-Soriano E, Fonseca P, Carneiro M, Moreira W. Seasonal variation in the chemical composition of two tropical seaweeds. Bioresource technology. 2006;97(18):2402-6.
9. Fujiwara-Arasaki T, Mino N, Kuroda M. The protein value in human nutrition of edible marine algae in Japan. Hydrobiologia. 1984;116(1):513-6.
10. Fleurence J. Seaweed proteins: biochemical, nutritional aspects and potential uses. Trends in Food Science & Technology. 1999;10(1):25-8.
11. Xiu-geng F, Ying B, Shan L. Seaweed cultivation: traditional way and its reformation. Chinese Journal of Oceanology and Limnology. 1999;17(3):193-9.

12. Mabeau S, Fleurence J. Seaweed in food products: biochemical and nutritional aspects. *Trends in Food Science & Technology*. 1993;4(4):103-7.
13. Patarra R, Leite J, Pereira R, Baptista J, Neto A. Fatty acid composition of selected macrophytes. 2012.
14. Bocanegra A, Bastida S, Bened íJ, R ódenas S, S ánchez-Muniz FJ. Characteristics and nutritional and cardiovascular-health properties of seaweeds. *Journal of medicinal food*. 2009;12(2):236-58.
15. Fern ández-Reiriz M, Perez-Camacho A, Ferreiro M, Blanco J, Planas M, Campos M, et al. Biomass production and variation in the biochemical profile (total protein, carbohydrates, RNA, lipids and fatty acids) of seven species of marine microalgae. *Aquaculture*. 1989;83(1):17-37.
16. Pomin VH, Mour ão PAS. Structure, biology, evolution, and medical importance of sulfated fucans and galactans. *Glycobiology*. 2008;18(12):1016-27.
17. Berteau O, Mulloy B. Sulfated fucans, fresh perspectives: structures, functions, and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology*. 2003;13(6):29R-40R.
18. Ruperez P. Mineral content of edible marine seaweeds. *Food Chemistry*. 2002;79(1):23-6.
19. McDermid KJ, Stuercke B. Nutritional composition of edible Hawaiian seaweeds. *Journal of Applied Phycology*. 2003;15(6):513-24.
20. Kusaykin M, Bakunina I, Sova V, Ermakova S, Kuznetsova T, Besednova N, et al. Structure, biological activity, and enzymatic transformation of fucoidans from the brown seaweeds. *Biotechnology journal*. 2008;3(7):904-15.
21. Li B, Lu F, Wei X, Zhao R. Fucoidan: structure and bioactivity. *Molecules*. 2008;13(8):1671-95.

22. Patel S. Therapeutic importance of sulfated polysaccharides from seaweeds: updating the recent findings. *3 Biotech.* 2012;1-15.
23. Silva T, Alves L, Queiroz KCS, Santos M, Marques C, Chavante S, et al. Partial characterization and anti-coagulant activity of a heterofucan from the brown seaweed *Padina gymnospora*. *Brazilian journal of medical and biological research.* 2005;38(4):523-33.
24. Pereira MS, Mulloy B, Mourão PAS. Structure and anti-coagulant activity of sulfated fucans. *Journal of Biological Chemistry.* 1999;274(12):7656-67.
25. Sithranga Boopathy N, Kathiresan K. Anticancer drugs from marine flora: an overview. *Journal of Oncology.* 2011;2010.
26. Ghosh T, Chattopadhyay K, Marschall M, Karmakar P, Mandal P, Ray B. Focus on antivirally active sulfated polysaccharides: from structure-activity analysis to clinical evaluation. *Glycobiology.* 2009;19(1):2-15.
27. Teas J, Irhimeh M. Dietary algae and HIV/AIDS: proof of concept clinical data. *Journal of Applied Phycology.* 2012:1-8.
28. Cumashi A, Ushakova NA, Preobrazhenskaya ME, D'Incecco A, Piccoli A, Totani L, et al. A comparative study of the anti-inflammatory, anti-coagulant, antiangiogenic, and antiadhesive activities of nine different fucoidans from brown seaweeds. *Glycobiology.* 2007;17(5):541-52.
29. Cho ML, Lee BY, You SG. Relationship between oversulfation and conformation of low and high molecular weight fucoidans and evaluation of their in vitro anticancer activity. *Molecules.* 2010;16(1):291-7.
30. Chung HJ, Jeun J, Hwang SJ, Jun HJ, Kweon DK, Lee SJ. Toxicological evaluation of fucoidan from *Undaria pinnatifida* in vitro and in vivo. *Phytotherapy Research.* 2010;24(7):1078-83.

31. Kim KJ, Lee OH, Lee HH, Lee BY. A 4-week repeated oral dose toxicity study of fucoidan from the *Sporophyll* of *Undaria pinnatifida* in Sprague–Dawley rats. *Toxicology*. 2010;267(1):154-8.
32. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes research and clinical practice*. 2011.
33. Harris MI. Classification, diagnostic criteria, and screening for diabetes. *Diabetes in America*. 1995;2:15-36.
34. Hanefeld M. Cardiovascular benefits and safety profile of acarbose therapy in prediabetes and established type 2 diabetes. *Cardiovascular diabetology*. 2007;6(1):20.
35. Atlas D. International Diabetes Federation. IDF-2000. 2000.
36. Lee SH, Karadeniz F, Kim MM, Kim SK. α - Glucosidase and α - amylase inhibitory activities of phloroglucinal derivatives from edible marine brown alga, *Ecklonia cava*. *Journal of the Science of Food and Agriculture*. 2009;89(9):1552-8.
37. Loizzo MR, Saab AM, Tundis R, Menichini F, Bonesi M, Piccolo V, et al. *In vitro* inhibitory activities of plants used in Lebanon traditional medicine against angiotensin converting enzyme (ACE) and digestive enzymes related to diabetes. *Journal of ethnopharmacology*. 2008;119(1):109-16.
38. Porte Jr D. Clinical importance of insulin secretion and its interaction with insulin resistance in the treatment of type 2 diabetes mellitus and its complications. *Diabetes/metabolism research and reviews*. 2001;17(3):181-8.
39. Lebovitz HE. Alpha-glucosidase inhibitors. *Endocrinology & Metabolism Clinics of North America*. 1997;26(3):539-51.
40. Inzucchi SE. Oral antihyperglycemic therapy for type 2 diabetes. *JAMA: the journal of the American Medical Association*. 2002;287(3):360-72.

41. Qiao Q, Larsen S, Borch-Johnsen K, Nissinen A, Pekkanen J, Tuomilehto J, et al. Glucose tolerance and cardiovascular mortality. Comparison of fasting and 2-hour diagnostic criteria. *Archives of Internal Medicine*. 2001;161(3):397-405.
42. Frantz S, Calvillo L, Tillmanns J, Elbing I, Dienesch C, Bischoff H, et al. Repetitive postprandial hyperglycemia increases cardiac ischemia/reperfusion injury: prevention by the α -glucosidase inhibitor acarbose. *The FASEB journal*. 2005;19(6):591-3.
43. Chiasson JL, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet*. 2002;359(9323):2072.
44. Holman RR, Cull CA, Turner RC. A randomized double-blind trial of acarbose in type 2 diabetes shows improved glycemic control over 3 years (UK Prospective Diabetes Study 44). *Diabetes Care*. 1999;22(6):960-4.
45. Chiasson JL, Josse RG, Leiter LA, Mihic M, Nathan DM, Palmason C, et al. The effect of acarbose on insulin sensitivity in subjects with impaired glucose tolerance. *Diabetes Care*. 1996;19(11):1190-3.
46. Dahanukar S, Kulkarni R, Rege N. Pharmacology of medicinal plants and natural products. *Indian Journal of Pharmacology*. 2000;32(4):81-118.
47. Cho M, Han JH, You S. Inhibitory effects of fucan sulfates on enzymatic hydrolysis of starch. *LWT-Food Science and Technology*. 2011;44(4):1164-71.

Chapter 2 Manuscript

***In vitro* investigation of inhibitory effect of fucoidan on α -glucosidase enzyme activity**

Xin Yin, Rahul Patel, Gregory Peterson, Uttam Budhathoki

School of Pharmacy, University of Tasmania, Sandy Bay, Hobart, Tasmania, Australia 7005

Corresponding author

Rahul Patel (Email Rahul.Patel@utas.edu.au)

Key words: fucoidan, acarbose, α -glucosidase, inhibitory assay

Abstract

Context. Diabetes causes a significant burden to the society and individuals. Current treatments exhibit many limitations. Therefore more effective and safer medications are required for its treatment.

Objective. The aim of this study was to investigate the inhibitory effects of fucoidans from *Undaria pinnatifida*, *Fucus vesiculosus* and *Fucus synergy* on the activity of α -glucosidase.

Materials and methods. Enzyme activity and enzyme kinetics of α -glucosidase enzyme extracted from rat-intestinal acetone powder were carried out using standard calibration curve of p-nitrophenol. Then enzyme inhibition assay was carried out using fucoidan from three different sources. Water and acarbose were used as negative and positive controls respectively.

Results. The enzyme activity, K_m and V_{max} of α -glucosidase were 0.0381 U ml^{-1} , $8.8339 \mu\text{g}$ and $0.2885 \mu\text{g ml}^{-1}$ respectively. Acarbose showed inhibition of α -glucosidase activity while three different types of tested fucoidan showed very little or no activity.

Discussion and conclusion. Experimental results showed negligible inhibitory effect of fucoidan on the activity of α -glucosidase.

Introduction

Diabetes is a group of heterogeneously metabolic disorders, characterised by hyperglycemia and glucose intolerance. It is mainly divided into insulin-dependent diabetes mellitus (IDDM) and non-insulin-dependent diabetes mellitus (NIDDM). Destruction of pancreatic beta-cells contributes to type I diabetes (IDDM), whereas insulin resistance or relative insulin deficiency is believed to be one major cause of type 2 diabetes (NIDDM) (1). Long-term abnormal glucose metabolism in diabetes provokes damage to blood vessels and double the risk of cardiovascular diseases (CVDs). The diabetic complications could be both macrovascular and microvascular, such as ischemic heart disease and blindness (2). Generally, the elderly patients with long-term poor blood glucose control are more likely to develop nephropathy, retinopathy, neuropathy and CVDs. By the end of 2000, 3% of the Western population had non-insulin-dependent diabetic mellitus (NIDDM) and 40% of these were patients over 60 years old. (3). Among these complications, CVDs such as coronary heart disease, stroke and peripheral vascular disease are the main reasons of mortality in diabetic patients. Many studies have indicated that postprandial hyperglycemia together with other risk factors, for example, smoking contribute to the development of CVDs (4). Therefore, preventing the onset of CVDs and correcting the postprandial hyperglycemic state are two core strategies in diabetic treatment. Current approaches include lifestyle modification and medication intervention (5). However, current pharmacological therapy is still facing a great challenge because evidence shows that some conventional oral anti-hyperglycemic agents increase the cardiovascular mortality (6). In the recent years, natural

products (7, 8) and chemically synthesised agents targeted on digestive starch enzymes, for example, α -glucosidase, have proven to be effective in treating postprandial hyperglycemia. Acarbose, a synthesised α -glucosidase inhibitor exhibited promising antidiabetic effect. It not only reduces postprandial hyperglycemia but also reduces the risk of CVDs by 49% (9). However, a study in 1994 revealed that approximately 46% of subjects complained about gastrointestinal discomfort such as flatulence and diarrhoea and about 2% did not tolerate acarbose (5). To reduce the side effects associated with synthesised α -glucosidase inhibitors, natural products extracted from marine algae such as fucoidan have been investigated for their potentially inhibitory effect on α -glucosidase.

Marine algae represents a large family of organisms and contains high content of sulfated polysaccharides which exhibit a variety of biological activities such as anti-oxidant (10), anti-viral and anti-bacterial (11), anti-tumor, anti-inflammatory and anti-coagulant (12). Fucoidan is a water soluble sulfated polysaccharide (13). A small in vitro study has indicated that fucoidan can potentially inhibit the activity of α -glucosidase enzyme (14) probably because of structural similarity to acarbose.

Many factors, such as the structure of fucoidan and types of seaweed sources can influence its inhibitory effect on α -glucosidase activity. Some studies also indicate that those factors make it hard to determine the anti-diabetic effect of fucoidan (15, 16). Hence, more research on anti-hyperglycemic activity of fucoidan is still needed. The objective of this study was to investigate the inhibitory effects of fucoidans from *Undaria pinnatifida*, *Fucus vesiculosus* and *Fucus synergy* on the activity of α -glucosidase.

Methods

Materials

Rat-intestinal acetone powder, sodium carbonate, 4-nitrophenol, acarbose, monobasic potassium phosphate, dibasic potassium phosphate and 4-nitrophenyl α -D-glucopyranoside (PNPG) were purchased from Sigma-Aldrich (New South Wales, Australia). Three different types of fucoidan (*Undaria pinnatifida*, *Fucus vesiculosus*, *Fucus synergy*) were generously donated by Marinova Pty Ltd (Tasmania, Australia).

Extraction of mammalian α -glucosidase

The method described by Mohamed and co-workers was modified for the extraction of α -glucosidase from rat-intestinal acetone powder (17). Briefly, rat-intestinal acetone powder (200 mg) was mixed with ice cold phosphate buffer (4 ml, 50 mM, pH 6.8). The suspension was then sonicated for 45 minutes using ultrasonic bath (Sonorex RK-100, Bendelin Electronic, Berlin, Germany) kept at approximately 0 to 4°C. The resultant mixture was centrifuged (Eppendorf centrifuge, model number 5417R, Hamburg, Germany) at 4°C and 13,000 rpm for 30 minutes. The supernatant was then carefully removed and stored at -20°C for further experiments.

Determination of enzyme activity

The activity of α -glucosidase was determined using PNPG (an enzyme substrate). The reaction between the enzyme and its substrate was initiated by mixing 10 μ l of enzyme extract and 20 μ l of 1 mM PNPG with 50 μ l of 50 mM phosphate buffer pH 6.8. The

enzyme–substrate mixture was then incubated in a Bioer mixing block (MB–102, Hangzhou, China) at 37 °C (500 rpm) for 30 minutes. The reaction was stopped using 50 µl of 0.1 M sodium carbonate. The intensity of colour, which was proportional to the activity of enzyme, was read at 405 nm using a micro–plate reader (Thermo Scientific Multiskan60, Massachusetts, United States of America). A solution of 4-nitrophenol (100 µg ml⁻¹) was diluted with a 50 mM of phosphate buffer to obtain 5 different concentrations (5, 10, 15, 20 and 25 µg ml⁻¹). A standard calibration curve (n=3) was generated by analysing these standards at 405 nm using a microtiter plate reader. Control samples (n=3) were prepared and analysed in a similar way except the enzyme was denatured by placing it in boiling water for 10 minute. The activity of enzyme (U ml⁻¹) was calculated using a following formula:

$$\text{enzyme activity} = \frac{m(\text{product}) \div \text{MW}(\text{product})}{30 \text{ mins} \times 0.01 \text{ ml}}$$

where m (product) is the mass of p–nitrophenol, MW (product) is the molecular weight of p–nitrophenol.

Enzyme kinetics of α–glucosidase

Kinetic study of the enzymatic reaction was carried out using different concentrations of PNPG in the presence of enzyme. A solution (n=3) containing 50µl of 50 mM phosphate buffer (pH 6.8), 10 µl of enzyme and 20 µl of different concentrations (ranging from 0.5 to 16 µg ml⁻¹) of PNPG was incubated at 37 °C using a mixing block for 30 minutes. The reaction was then stopped by adding 50 µl of 0.1 M sodium carbonate, followed by incubation in mixing block for approximately 1 minute. The intensity of colour was measured

at 405 nm using a microtiter plate reader. Control samples (n=3) were prepared and analysed in a similar way except the enzyme was denatured by placing it in boiling water for 10 minutes. Samples were prepared in triplicate and analysed in duplicate. Lineweaver–Burk plot was used to determine maximum reaction rate (V_{\max}) and Michaelis constant (K_m) of the enzyme (17, 18).

Inhibitory assay

The inhibitory effect of fucoidan on α -glucosidase was investigated using varying concentrations (ranging from 0.01 to 1000 $\mu\text{g/ml}$) of three different types of fucoidan (*Undaria Pinnatifida*, *Fucus Vesculosus*, *Fucus Synergy*). Enzyme (0.04U ml^{-1} , 50 μl) was mixed with 20 μl of each concentration of inhibitor in eppendorf tube and incubated for 10 minutes at 37 °C and 500 rpm. Technically, various concentrations of inhibitor were diluted from a stock solution (7 mg ml^{-1}) which provided 1000 $\mu\text{g/ml}$ of fucoidan or acarbose in the final reaction mixture (140 μl). The fucoidan powder (70 mg) was placed in a 10 ml volumetric flask and then was sonicated using ultrasonic bath until all particles dissolved. After this, the stock solution was diluted to obtain different concentrations (0.05, 0.1, 1, 2.5, 7.5, 10, 50, 100, 500, 1000 $\mu\text{g ml}^{-1}$) of fucoidan. The preparation of positive control (acarbose) followed a similar pattern, whereas Milli-Q water was used instead of tested sample in the negative controls. Enzyme (0.04U/ml) and its activity were obtained based on the process of extraction of mammalian α -glucosidase enzyme and determination of enzyme activity respectively. A solution (n=3) containing enzyme extract and different concentrations of fucoidan was incubated at 37°C for 10 minutes. PNPG (1 mM, 20 μl) was added to the

enzyme–fucoidan mixture to initiate the enzyme–substrate reaction. The mixture was then incubated for 30 minutes at 37°C in a mixing block. Sodium bicarbonate (50 µl of 0.1M) was added to stop the enzyme–substrate reaction. At end of assay, the intensity of the colour was measured at 405 nm using a microtiter plate reader. The inhibitory effect of fucoidan and positive control were calculated using a following formula:

$$\% \text{ inhibition} = \frac{\text{MC} - \text{MS}}{\text{MC}} \times 100$$

where MC and MS are mass of product converted in the negative control and that of sample or the positive control respectively.

Results

Enzyme activity of α -glucosidase

Enzyme activities of four different extracts were found to be 0.0076, 0.0098, 0.0237 and 0.0381 U ml⁻¹. The fourth enzyme extract (0.0381 U ml⁻¹) was used for the enzyme inhibitory assay.

Enzyme inhibitory assay of α -glucosidase

Three different extracts of fucoidan assessed in enzyme inhibitory assay had negligible effect when compared with acarbose. The maximum percentage of inhibition of acarbose was 61.11%, achieved at a concentration of 1000 μ g ml⁻¹, and showed a consistent tendency during the enzyme inhibitory assay. In contrast, the maximum percentage of inhibition of fucoidan extracted from *Fucus Synergy* was 7.57%, followed by the extracts of *Fucus vesiculosus* (6.14%) and *Undaria pinnatifida* (3.30%) (Figure 1). As fucoidan did not show increase in inhibition with increase in concentration of it, it can be confirmed that all the tested fucoidan extracts did not show inhibitory effect on the enzyme.

V_{max} and K_m of α -glucosidase

The enzyme kinetics study showed that the enzyme had K_m and V_{max} values of 8.8339 μ g ml⁻¹ and 0.2885 μ g min⁻¹ respectively (Figure 2).

Discussion

Although the inhibitory effect of fucoidan from two commercial seaweeds (*Fucus vesiculosus* and *Ascophyllum nodosum*) on yeast α -glucosidase has been reported in 2012 (19), this study re-assessed its anti-hyperglycemic effect on the same enzyme for three main reasons. First of all, yeast α -glucosidase instead of mammalian source was used in previous study (19). The second reason to revisit is they have used four concentrations of fucoidan ranging from 0.005 to 0.05 $\mu\text{g ml}^{-1}$ in 5 ml of final enzyme–substrate–inhibitor solution where as higher concentration of fucoidan is included in my experiment. Moreover they used negative control only but both negative and positive controls were used in our study. Third reason is that the activity of fucoidan was not investigated in duplicate or triplicate. Multiple analysis of the same sample is important for the reliability of the results (20).

Another study in 2012 showed that fucoidan extract from *Fucus vesiculosus* showed the maximum percentage inhibition of 51.4% (19), whereas fucoidan from same source and other two sources did not exhibit inhibition on the same enzyme in our study. This could be due to property of fucoidan shows highly species–specific characteristics and some other parameters such as harvesting period and growth conditions also contribute to variance on its inhibitory effect (13).

The extraction of enzyme was influenced by sonication time. Increase in sonication time increased the amount of enzyme that can be extracted from the rat intestinal powder, thus higher enzyme activity was obtained with longer sonication time. For example, the enzyme

activities of α -glucosidase was 0.0076 U ml^{-1} after 30 minutes of sonication time, compared to 0.0381 U ml^{-1} after 60 minutes of sonication.

Michaelis constant (K_m) in enzyme kinetics indicated the affinity of α -glucosidase to PNPG. K_m also equals to the concentration of substrate at which the reaction takes place at one half of its maximum reaction rate and this value is reversely proportional to the affinity (21). In this study, K_m value indicated the required concentration of PNPG and the reaction time to achieve equilibrium. Relevant studies on mammalian α -glucosidase are quite limited. However, one study in 1983 showed K_m and V_{max} values of yeast α -glucosidase from *Lactobacillus acidophilus* on PNPG as $7.9988 \text{ } \mu\text{g ml}^{-1}$ and $0.001807 \text{ } \mu\text{g min}^{-1}$ respectively after 30 minutes reaction (22). My experimental result showed that K_m value was similar to yeast α -glucosidase but V_{max} was 160 times higher than it. That means yeast α -glucosidase and mammalian α -glucosidase have similar affinity to PNPG, whereas the rate of transformation from substrate to product by using rat α -glucosidase is 160 times more than that of yeast α -glucosidase. Additionally, according to the K_m value, the concentration of PNPG at which the reaction takes place at half of V_{max} is $0.0635 \text{ } \mu\text{M}$; therefore, only approximate $0.127 \text{ } \mu\text{M}$ of PNPG is required to achieve the V_{max} , compared with 1 mM of PNPG employed in this study.

However, due to time constrain it was not possible to investigate the activity of fucoidan extracted from other sources. Also, this study did not investigate the influence of various parameters such as seasonal variation and extraction procedures on the inhibitory effect of fucoidan.

Conclusions

The inhibitory effects of three types of fucoidan were negligible, compare with acarbose, which reached a maximum percentage inhibition of 61% at 30 minutes reaction time using $1000\text{ }\mu\text{g ml}^{-1}$ acarbose. The V_{max} and K_m values of rat α -glucosidase were $0.2885\text{ }\mu\text{g min}^{-1}$ and $8.8339\text{ }\mu\text{g ml}^{-1}$ respectively. Enzyme activity of α -glucosidase was 0.0381 U ml^{-1} . According to the discussion above, α -glucosidase has high affinity to PNPG. The concentration of PNPG (1 mM) and the reaction time (30 minutes) were sufficient to achieve the V_{max} .

Declaration of interest

The author has declared no potential conflicts of interest.

List of figures

Figure 1

The inhibitory effect of fucoidan extracted from *Fucus synergy*, *Undaria pinnatifida* and *Fucus vesiculosus* on the activity of α -glucosidase, in comparison with acarbose at the same concentrations





Figure 2

Partial determination of enzyme kinetics of α -glucosidase using different concentrations of PNPG (ranging from 0.5 to 16 $\mu\text{g ml}^{-1}$). The reciprocal values of intercept of X axis and Y axis indicate V_{max} and K_m respectively.



References

1. Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabetic medicine*. 1997;14(S5):S7-S85.
2. Costantino L, Rastelli G, Vianello P, Cignarella G, Barlocco D. Diabetes complications and their potential prevention: aldose reductase inhibition and other approaches. *Medicinal research reviews*. 2000;19(1):3-23.
3. Bonora E, Muggeo M. Postprandial blood glucose as a risk factor for cardiovascular disease in type II diabetes: the epidemiological evidence. *Diabetologia*. 2001;44(12):2107-14.
4. Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care*. 1993;16(2):434-44.
5. Hanefeld M. Cardiovascular benefits and safety profile of acarbose therapy in prediabetes and established type 2 diabetes. *Cardiovascular diabetology*. 2007;6(1):20.
6. Laakso M. Hyperglycemia and cardiovascular disease in type 2 diabetes. *Diabetes*. 1999;48(5):937-42.
7. Jung M, Park M, Lee HC, Kang Y-H, Kang ES, Kim SK. Antidiabetic agents from medicinal plants. *Current Medicinal Chemistry*. 2006;13(10):1203-18.
8. Kumar S, Narwal S, Kumar V, Prakash O. α -glucosidase inhibitors from plants: A natural approach to treat diabetes. *Pharmacognosy reviews*. 2011;5(9):19.

9. Ceriello A. Postprandial Hyperglycemia and Diabetes Complications Is It Time to Treat? *Diabetes*. 2005;54(1):1-7.
10. Rupérez P, Ahrazem O, Leal JA. Potential anti-oxidant capacity of sulfated polysaccharides from the edible marine brown seaweed *Fucus vesiculosus*. *Journal of agricultural and food chemistry*. 2002;50(4):840-5.
11. Mayer A, Hamann MT. Marine pharmacology in 2001–2002: marine compounds with anthelmintic, antibacterial, anti-coagulant, antidiabetic, antifungal, anti-inflammatory, antimalarial, antiplatelet, antiprotozoal, antituberculosis, and antiviral activities; affecting the cardiovascular, immune and nervous systems and other miscellaneous mechanisms of action. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2005;140(3):265-86.
12. Cumashi A, Ushakova NA, Preobrazhenskaya ME, D'Incecco A, Piccoli A, Totani L, et al. A comparative study of the anti-inflammatory, anti-coagulant, antiangiogenic, and antiadhesive activities of nine different fucoidans from brown seaweeds. *Glycobiology*. 2007;17(5):541-52.
13. Berteau O, Mulloy B. Sulfated fucans, fresh perspectives: structures, functions, and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology*. 2003;13(6):29R-40R.
14. Apostolidis E, Lee C. In Vitro Potential of *Ascophyllum nodosum* Phenolic Anti-oxidant - Mediated α - Glucosidase and α - Amylase Inhibition. *Journal of food science*. 2010;75(3):H97-H102.

15. Ale MT, Mikkelsen JD, Meyer AS. Important determinants for fucoidan bioactivity: A critical review of structure-function relations and extraction methods for fucose-containing sulfated polysaccharides from brown seaweeds. *Marine drugs*. 2011;9(10):2106-30.
16. Li B, LU F, SUN K-x. Research progress on bioactivity of fucoidan. *FOOD AND MACHINERY*. 2006;22(3):157.
17. Shihabudeen HMS, Priscilla DH, Thirumurugan K. Cinnamon extract inhibits α -glucosidase activity and dampens postprandial glucose excursion in diabetic rats. 2011.
18. Kim YM, Jeong YK, Wang MH, Lee WY, Rhee HI. Inhibitory effect of pine extract on α -glucosidase activity and postprandial hyperglycemia. *Nutrition*. 2005;21(6):756-61.
19. KIM KTAE. Seasonal variation of seaweed components and novel biological function of fucoidan extracted from brown algae in Quebec. 2012.
20. Mason RL, Gunst RF, Hess JL. Statistical design and analysis of experiments: with applications to engineering and science: Wiley-Interscience; 2003.
21. Cornish-Bowden A. Fundamentals of enzyme kinetics: Butterworths London; 1979.
22. Li K, Chan K. Production and properties of alpha-glucosidase from *Lactobacillus acidophilus*. *Applied and environmental microbiology*. 1983;46(6):1380-7.